

Africanization of Melliferous Bees (*Apis mellifera*.L.) Bibliographic Review

Diego Masaquiza Moposita¹, Lino Miguel Curbelo Rodríguez² & Amilcar Arenal Cruz³

¹ORCID <https://orcid.org/0000-0001-5176-8261>, Ignacio Agramonte Loynaz University of Camagüey, Center for Animal Production Studies, Camagüey, Cuba, ²ORCID <https://orcid.org/0000-0003-0453-2357>, Ignacio Agramonte Loynaz University of Camagüey, Center for Animal Production Studies, Camagüey, Cuba, ³ORCID <https://orcid.org/0000-0003-2912-9871>, Ignacio Agramonte Loynaz University of Camagüey, Center of Molecular Biology, Camagüey, Cuba.

Citation: Masaquiza Moposita, D., Curbelo Rodríguez, L., & Arenal Cruz, A. (2020). Africanization of Melliferous Bees (*Apis mellifera*.L.) Bibliographic Review. *Agrisost*, 26(2), 1-12. <https://doi.org/10.5281/zenodo.7573827>

Received: February 2, 2020

Accepted: February 22, 2020

Published: March 2, 2020

Funding source: Not declared.

Conflicts of interest: No conflict of interest has been declared.

Email: diegomasaquiza@reduc.edu.cu

Abstract

Context: The hybridization process (Africanization) of European bees with African bees is a problem for apiculture farmers in the Americas, due to high swarming levels and defensiveness. The latter hinders colony handling, and has caused accidents to people and animals, increasing the risks of beekeeping. In this sense, there is a need for genetic breeding of melliferous bees, and previous identification of subspecies.

Aim: To evaluate the origin of melliferous bees (*Apis mellifera*), and the process of Africanization and dispersion of Africanized bees throughout the Americas, as well as methods of identification.

Methods: The databases of Scienedirect, Google-Scholar, Scopus, and NCBI were reviewed under the following key words, *Apis mellifera*, *Apis*, Africanized bees, geometric morphometrics, mitochondrial DNA. Special emphasis was paid to papers published within the last five years.

Results: The origin and distribution of melliferous bees, and the Africanization and dispersion processes of Africanized bees were described. Additionally, the evolution of methods for the characterization of *Apis mellifera* species were updated.

Conclusions: Africanization can be considered the most important process in the transformation of conduct and morphological features of melliferous bees, which allowed for their rapid dispersion in the Americas. The identification methods based on parents are essential to know possible process of genetic erosion, and to present strategies for bee conservation and breeding in every region.

Key words: *mellifera*, *Apis*, Africanized bees, geometric morphometrics, mitochondrial DNA.

Introduction

Africanization is one of the fastest and most striking biological invasions ever known worldwide (Branchiccela et al., 2014; Medina-Flores et al., 2015), and there are divergent criteria in favor and against the results for apiculture in the Americas. Several actions are being taken to control this process, such as the introduction of European germplasm, and colony isolation, though neither has stopped these effects. Accordingly, learning to live with these bees seems to be the logical solution, since these species have positive traits, such as a marked hygienic behavior, high production of honey, and so on. Then, it is important to characterize bees to learn the existing genetic material in every region, and set actions in the mid and long terms, in order to reduce defensiveness and swarming, which are negative

traits found in the Africanized hybrids. The rationale for these arguments aims to examine the origin of melliferous bees (*Apis mellifera*), the process of dispersion of Africanized bees through the Americas, and the methods of bee identification.

Development

Origin and distribution of *Apis mellifera*

The evolution of class *Insecta*, and order *Hymenoptera* took place in the Jurassic period, approximately 180 millions of years ago, as a result of the co-evolution of flower plants, which had ovaries and stigmas, and depended on insects for entomophile reproduction-pollinizing (Carpana, 2004). The diversity of species is the result from radiation in the early Pleistocene period due to

climatic change and habitats during the last ice age (Combey, Quindahor & Mensah, 2018).

Of the three subfamilies of the Apidae family (Meliponinae, Bombinae, and Apinae), Apinae withstood the best adaptation, which made them cosmopolitan (Verde, 2014). *A. mellifera* is the only bee species that evolved in Europe, the Middle East, and Africa, where several subspecies can be identified (Genchi, Reynaldi & Bravi, 2018). Depending on the geographical origin, there are 29 subspecies (Valido, Rodríguez-Rodríguez & Jordano, 2014; Abizanda, 2018) with remarkable morphological and genetic differences (Wallberg et al., 2014), with the largest distribution worldwide, and tropical and temperate ecotypes.

Subspecies are grouped in five evolutionary lines, mainly determined by the climatic and flora variations, as a result of glacier and post-glacier periods (Miguel et al., 2016).

Meixner et al. (2013) noted that line A includes African subspecies *A. m. intermissa* and *A. m. sahariensis* in the North; *A. m. lamarckii*, *A. m. simensis*, and *A. m. adansonii* in the west and center; *A. m. scutellata* in the center and south; *A. m. capensis* in the south, *A. m. monticola* in the mountainous region of southeastern Africa, *A. m. litorea* in the southeast of Africa, and finally *A. m. unicolor* which is endemic to Madagascar (Techer et al., 2017). Line M is made of Western Europe subspecies, including *A. m. mellifera*; *A. m. iberiensis* (Kandemir, Özkan & Fuchs, 2011). Line C is made of Eastern Europe subspecies, including *A. m. ligustica*, *A. m. carnica*. Line O includes the Middle East subspecies (Rúa, Martínez, Domingo & Gabaldón, 2013; Charistos, Hatjina, Bouga, Mladenovic & Maistros, 2014; Wallberg et al., 2014). And Y includes subspecies *A. m. jemenitica* from Ethiopia (Meixner et al., 2013). Line Z, with populations spread in Libya (Péntek-Zakar, Oleksa, Borowik & Kusza, 2015)

In the sixteenth century, the European settlers introduced melliferous bees (*A. mellifera*) in the Americas; the data show the dispersion process in North America in 1622, Cuba, 1763, and Brazil and Chile 1839 (Nogueira-Neto, 1972). In Ecuador, the first bee-hives were introduced in 1870 with the honey producing European ecotype *A. m. ligustica* (Hidalgo & Mena, 2003).

In the second half of the twentieth century, in most of the Americas, a biological invasion known as Africanization took place (Kerr, 1967). This phenomenon (Garcia et al., 2013) resulted from the escape of African bees (*Apis mellifera scutellata*) from an experimental beekeeping facility in Brazil, in 1957.

The Africanization of melliferous bees in 1956, *A. m. scutellata*, African line, was introduced in Brazil from South Africa and Tanzania (Chapman et al., 2015), which started as a breeding project to increase honey production of European bees in the tropical climate of the Amazon basin. The European bees are unable to link their biological cycle to the duration of the day in the tropic. However, flowering in temperate areas is framed, and in the tropics, flowering responds to rainy periods (Moore, Wilson & Skinner, 2015).

The idea was to create a “super bee”, a hybrid between European and African subspecies with proper temperament and search habits (Whitfield et al., 2006).

Nevertheless, Africanized bees have a high genetic variability due to their hybrid character; they are also known by the high productivity and tolerance against pathogens and parasites like *Varroa destructor* (Strauss, Dietemann, Human, Crewe & Pirk, 2015; Tibatá et al., 2018).

Factors like genetic plasticity and adaptability of bees (Sousa, Araújo, Gramacho & Nunes, 2016), and the wide geographical dispersion and environmental variability in the Americas, allow for hybrid evolutionary processes (Nunes, Araújo, Marchini & Moreti, 2012). By being in contact, the African and European colonies, the process of hybridization lead to the replacement and displacement of European traits by African ones (Rangel et al., 2016). Africanized bees are characterized by their high defensive behavior (DB), and migration (Medina-Flores et al., 2015). Their propagation causes problems to apiculture, and had important social and economic impacts due to their high defensiveness (Byatt, Chapman, Latty & Oldroyd, 2016).

Today, Africanized bees are present from the mid and south of the United States (Szalanski & Tripodi, 2014; Portman, Tepedino, Tripodi, Szalanski & Durham, 2018), Central America to the north of Argentina (Genchi et al., 2018), covering 20 million km² approximately. Their high colonizing activity is one of the fastest biological invasions known (Branchiccela et al., 2014; Medina-Flores et al., 2015).

Jarnevich et al. (2014) noted that hybrid Africanized bees spread out, and continue to do so in the continent (Figure 1). The first swarms of African bees arrived in Ecuador, Colombia, and Peru in the 1970s (Graciano, 2018).




Fig. 1. Dispersion of Africanized bees in the Americas (Winston, 1992).

Wallberg et al. (2014) claimed that Africanized bees in South America are mostly from African parents (70-90%), and in the United States (50-75%). This distribution probably shows the adaptation to local environmental conditions, and hybridization between melliferous bee subspecies (Coroian et al., 2014). The Africanization level may have implications in adaptability, biological attitude, behavior, pollinizing, and production of honey by *A. mellifera* (Mendoza et al., 2014; Hamiduzzaman et al., 2015). In this sense, the conservation of the genetic diversity of melliferous bees is important to preserve the high genetic adaptability of populations in different zones (Büchler et al., 2014), which is closely related to them.

Evolution of characterization methods of subspecies *Apis mellifera*

Since many years ago, several methods are being used to identify and classify *A. mellifera*, one of them is traditional morphometrics (Alpatov, 1929; Ruttner, 1988). Moreover, quite a few methods for morphological and molecular analysis are used for identification of *A. mellifera* hybrid, such as the Fast Africanized Bee Identification System (FABIS). This method permits preliminary identification of Africanized bees (Sylvester & Rinderer, 1987).

The Universal System to Detect Africanization (USDA-ID) used to declare official cases of Africanization (Sanford, 2006), is painstaking, since it requires twenty-five body parts or structures from each specimen. The Automated Bee Identification System (ABIS) that compares wing-vein graphs with a digital image from the anterior part of the sample, requires two minutes per sample, with an estimated

confidence of 98.05% between species, and 94% between subspecies of melliferous bees (Francoy et al., 2008).

Currently, geometric morphometrics methods have shown significant results in classification and identification of insects. This method provides a powerful tool to study the biological and phylogenetic classifications (Su, Cai & Huang, 2018). The geomorphometric methods can be used to solve taxonomy problems, and to visualize key differences in the shape of species (Canal, Hernández-Ortiz, Salas & Selivon, 2015; Qubaiová, Růžicka & Šípková, 2015).

The classification of different lines is supported by more accurate methods, such as biochemical methods: isoenzymes and aloenzymes (Nunamaker & Wilson, 1982; Ivanova et al., 2012); nuclear DNA molecular markers (Charistos et al., 2014); mitochondrial DNA (Szalanski & Tripodi, 2014); sequencing (Branchiccela et al., 2014; Kono & Kohn, 2015); microsatellites (Rangel et al., 2016), and cuticle hydrocarbons (Oyerinde, Salako & Rabiou, 2017). In recent years, the utilization of single nucleotide polymorphisms (SNP) has become relevant (Chávez-Galarza et al., 2015; Chapmaneta, 2015; Muñoz et al., 2015).

Morphometrics

The concept of beekeeping taxonomy was introduced by Dupraw (1965), and it was developed by Ruttner, Tassencourt & Louveaux (1978).

The first subspecific classifications were based on geographical and conduct features, depending on their distribution (Ruttner et al., 1978; Ruttner, 1988). Using morphometrics, diagnostic methods can be utilized in multiple measurements of many individuals; in physical and structural variations, which are universal features of a population, genus, species or subspecies (Darger, 2013). Now, subspecies differentiation is based on morphometric characteristics (Kandemir et al., 2011; Kono & Kohn, 2015; Sousa et al., 2016), which undergo gradual changes between stages, especially between subspecies that often just have a slight difference in the mean values (Abizanda, 2018).

Traditional morphometrics

The traditional morphometrics allowed for identification and classification of *A. mellifera* (Daly, Hoelmer, Norman & Allen, 1982; Ruttner, 1988). Variance and covariance studies of distance are used between point pairs, usually length and width of structures, in addition to proportions and angles (Rohlf & Marcus, 1993). The morphometric characteristics of bees are used, on the other hand, to identify breeds or geographical subspecies. Among

the first morphometric classification techniques are the ones established by Cochlov, in 1916, cited by Ruttner (1988), who compared proboscis length among six geographically different breeds of *A. mellifera*.

Later, new sets of characteristics were introduced by Goetze, in 1940, like hair length in the terga, the width of the first tarsus segment, and the length diameter of terga 3 and 4 (Ruttner, 1988), which were very efficient for *A. mellifera* subspecies discrimination. Howell, Daly & Balling (1978), developed the first efficient procedure, which consisted in measuring 25 characters by hand: wings, sternite, and legs, using an optical micrometer, with reliable results. These measurements were analyzed through multivariate statistics for identification. Later, Daly et al. (1982), improved this method with the inclusion of digital measurements (measurement system linked to a computer), that led to a much easier and faster procedure.

Ruttner (1988) perfected this technique, and described 36 morphometric features (distances, angles, pigmentation classes) for the discrimination of melliferous bees subspecies. These features helped establish four different evolutionary lines of melliferous bees (Ruttner, 1992).

The traditional morphometrics includes wing and body features that offer substantial understanding and proper discrimination among bee subspecies and populations. In the search for method improvements, 10 features described by Ruttner, in 1988 were used (Meixner et al., 2013).

The efficiency of this classification depends on the selection of relevant features that prevent mistaken information (da Silva et al., 2015). However, it is still used in various research to characterize subspecies in locations such as the mountain forest of Crimea (Bykova, Triseleva, Ivashov & Safonkin, 2016), in the Republic of Benin (Amakpe et al., 2018), Colombia (Graciano, 2018).

Geometric morphometrics

With the transition from descriptive to quantitative morphometrics, morphological identification is the most accurate and reproducible, since it relies on computerized techniques (Adams, Rohlf & Slice, 2004). The descriptive analysis of size and shape variations is a fundamental tool for organic biology studies (De Souza et al., 2015), which became popular in the late twentieth century, with the fusion of biology and geometry, in order to collect geometric information from biological forms, for comparative purposes (Torcida & Perez, 2012).

From the beginning, geometric morphology (GM) showed better descriptive power, in addition to an

approach that offers a relative description of the form by using reference points (Santana et al., 2014). It relies on integrated statistical analysis to collect spatial information of morphological structures, thus increasing accuracy in relation to traditional morphometry (Francoy, de Faria Franco & Roubik, 2012). It allows for specific analysis of morphometric variations of particular structure in organisms with different sizes (Nunes et al., 2012).

Geometric morphometry uses the coordinates of reference points, which are then overlapped through displacement, scale-up, and rotation, to remove the size effect (da Silva, Grassi, Sella, Francoy & Reali, 2015). Following overlapping, the landmark configurations only differ in their shape, and can be analyzed through multivariate statistical methods (Zelditch, Swiderski, Sheets & Fink, 2004).

Variation in the shape of the wing of bees at different taxonomic levels, with the utilization of GM is useful to discriminate subspecies (Charistos et al., 2014; De Souza et al., 2015; Oleksa & Tofilski, 2015; Abizanda, 2018), and the geographical variation of species (Hall, MacLeod & Wardhana, 2014; Carvajal et al., 2015). Furthermore, since the morphometric characteristics of bees are highly inherited, this technique can be suitable for contrasting.

GM was highly effective to discriminate 24 known subspecies of melliferous bees in Europe (Kandemir et al., 2011). In turn, Oleksa & Tofilski (2015) indicated that GM provides similar discrimination values to microsatellites. The high accuracy of classification indicates that the front wings carry sufficient information to distinguish the groups of bees examined (Charistos et al., 2014). Besides identifying hybridization process, the morphometric techniques are related to nuclear microsatellite analysis that measure changes in the size and shape of bee bodies, since they are controlled by nuclear genes (parent inheritance), in keeping with the arguments of Mortensen & Ellis (2015).

They also allow for optimization of analysis time by reducing the number of individuals used; 10-15 bees are recommended for morphometric analysis (Kandemir et al., 2011; Meixner et al., 2013; Sousa et al., 2016). The number of individuals may be reduced to five, to analyze more stable traits, such as wing veins (Francoy et al., 2008), with a high level of consistency between the wing and molecular information.

Molecular Techniques

The different molecular analyses performed to study the genetic diversity of *A. mellifera*, provide an accurate tool compared to classic morphometrics (Szalanski & Tripodi, 2014). Melliferous bee hybrids are morphologically similar, and hard to distinguish

from one another, so in some cases molecular analysis is necessary (Achou et al., 2015). In the last 35 years, the identification and distribution of different *A. mellifera* lines based on molecular and mitochondrial markers has been reported, (Wallberg et al., 2014) (Table 1)

Table 1. Recent molecular biology research to identify the maternal origin of melliferous bees

Analysis	Genes	Subspecies	Author
Analysis of polymorphism restriction	tRNAle u-COII, Cyt b, rRNA – COI, COI	<i>Apis mellifera</i>	Wu, Lu & Lu (2017)
Analysis of polymorphism s and sequencing	tRNAle u- COII	<i>Apis mellifera</i>	Branchicella et al. (2014)
Sequencing	mtDNA	<i>A. m. scutellata</i> <i>A. m. capensis</i>	Eimanifar, Kimball, Braun & Ellis (2018)
Sequencing	COI - COII	<i>Apis mellifera</i>	Magnus & Szalanski (2010)
Sequencing	COI - COII; ND2	<i>A. m. unicolor</i>	Techer et al. (2017)
Analysis of polymorphism restriction	COI - COII	<i>Apis mellifera</i>	Kelomey et al. (2017)
Analysis of polymorphism restriction	COI - COII	<i>A. m. intermissa</i> <i>A. m. sahariensis</i>	Achou et al. (2015)
PCR - RFLP	Cyt b	<i>A. m. scutellata</i>	Mortensen & Ellis (2015)
Sequencing	tRNAle u- COII	<i>Apis mellifera</i>	Szalanski & Magnus (2010)
Sequencing	COI - COII	<i>Apis mellifera</i>	Techer et al. (2015)
Sequencing	tRNAle u- COII	<i>A. m. iberiensis</i>	Chávez-Galarza et al. (2017)
Sequencing	mtDNA - Cyt-b	<i>Apis mellifera</i>	Szalanski & Tripodi (2014)

Mitochondrial DNA

The first molecular studies relied on the study of highly variable mitochondrial intergene COI-COII

locus (Péntek-Zakar et al., 2015); they were started in the early 1990s (Garnery, Solignac, Celebrano & Cornuet, 1993). It is a frequently used tool to discriminate evolutionary lines in melliferous bees populations (Rortais, Arnold, Alburaki, Legout & Garnery, 2011).

Mitochondrial DNA (mtDNA) is a tiny, 16 000 base pairs, circular molecule, which is completely transmitted by the queens to their offspring: workers and drones (Meixner et al., 2013). The mother inheritance of mtDNA in the bees is a molecular marker that allows for tracing the origin and genetic structure of melliferous bees. The study of a single individual provides direct information from the hive (Rúa et al., 2013), since this circular molecule is not changed by recombination, and it is transmitted without altering the offspring, except in cases of punctual mutations.

The mitochondrial genome is characterized by conserved gene contents, high level of nucleotide substitutions, and as a provider of a single parent marker (Péntek-Zakar et al., 2015). The mitochondrial chromosome of *A. mellifera* contains 13 genes that codify for proteins, two ribosomal genes, 22 transference RNA, and a region which is rich in A + T. It is conservative in terms of their position (Crozier & Crozier, 1993).

The intergene COI-COII locus is made of two distinct nucleotide sequences, named P and Q, where P can appear in several variations (P0, P1, P2). The variations observed in the sequences of the region can be used to contrast the five lines of melliferous bees, and to discriminate the subspecies of *A. mellifera* from one another (Alattal et al., 2014). According to Garnery et al. (1993), each evolutionary line includes a P sequence variant combined with one of the four copies of the Q sequence found in that locus, except for the C line subspecies, which do not have a copy of the P sequence, and only possess a Q sequence copy (Rúa et al., 2013).

The codifying insert of the intergene locus helps distinguish the African line from the European one, since the melliferous bees of the east European line (C) have a shorter intergene sequence due to the absence of P, and a single copy of Q (Meixner et al., 2013).

Melliferous bees that belong to lines M, A, Z, and Y have longer intergene regions, because they contain between one and five Q elements (Rortais et al., 2011), in addition of a variant of the P element.

The mitochondrial chromosome of *A. mellifera* (Figure 2) contains 13 genes that codify for proteins, two ribosomal genes, 22 transference RNA, and a locus that is rich in A + T. It is conservative in terms

Lino Miguel Curbelo Rodríguez: research planning, data collection, analysis and interpretation of results, redaction and final review of the manuscript.

Amílcar Arenal: research planning, data collection, analysis and interpretation of results, redaction and final review of the manuscript.

Conflicts of interest

No conflict of interest has been declared.

References

- Abizanda, C. (2018). *Caracterización morfológica y molecular de las abejas melíferas en la provincia de Huesca*. (Trabajo Fin de Máster), Universidad de Zaragoza, Escuela Politécnica Superior, España. Retrieved on March 12, 2019, from: <http://zaguan.unizar.es/record/69789/files/TAZ-TFM-2018-063.pdf?version=1>
- Achou, M., Loucif-Ayad, W., Legout, H., Hmidan, H., Alburaki, M., & Garnery, L. (2015). An insightful molecular analysis reveals foreign honey bees among Algerian honey bee populations (*Apis mellifera* L.). *Data Mining Genomics Proteomics*, 6(1), 166, doi: <https://doi.org/10.4172/2153-0602.1000166>
- Adams, D.C., Rohlf, F. J., & Slice, D. E. (2004). Geometric morphometrics: ten years of progress following the 'revolution'. *Italian Journal of Zoology*, 71(1), 5-16, doi: <https://doi.org/10.1080/11250000409356545>
- Alattal, Y., Alsharhi, M., Alghamdi, A., Alfaify, S., Migdadi, H., & Ansari, M. (2014). Characterization of the native honey bee subspecies in Saudi Arabia using the medina COI-COI intergenic region and morphometric characteristics. *Bulletin of Insectology*, 67(31-37). Retrieved on April 5, 2019, from: <https://pdfs.semanticscholar.org/0222/f2c52f135ab19ce51de8247129de23b6b58b.pdf>
- Alpatov, W. (1929). Biometrical studies on variation and races of the honey bee (*Apis mellifera* L.). *The Quarterly Review of Biology*, 4(1), 1-58, doi: <https://doi.org/10.1086/394322>
- Amakpe, F., De Smet, L., Brunain, M., Frans, J., Sinsin, B., & de Graaf, D. (2018). Characterization of native honey bee subspecies in Republic of Benin using morphometric and genetic tools. *J. Apic. Sci.*, 62(1), 47-59, doi: <https://doi.org/10.2478/jas-2018-0006>
- Branchiccela, B., Aguirre, C., Parra, G., Estay, P., Zunino, P., & Antúnez, K. (2014). Genetic changes in *Apis mellifera* after 40 years of Africanization. *Apidologie*, 45(6), 752-756, doi: <https://doi.org/10.1007/s13592-014-0293-2>
- Büchler, R., Costa, C., Hatjina, F., Andonov, S., Meixner, M. D., Le Conte, Y., . . . Wilde, J. (2014). The influence of genetic origin and its interaction with environmental effects on the survival of *Apis mellifera* L. colonies in Europe. *Journal of Apicultural Research*, 53(2), 205-214, doi: <https://doi.org/10.3896/IBRA.1.53.2.03>
- Byatt, M. A., Chapman, N. C., Latty, T., & Oldroyd, B. P. (2016). The genetic consequences of the anthropogenic movement of social bees. *Insectes Sociaux*, 63(1), 15-24, doi: <https://doi.org/10.1007/s00040-015-0441-3>
- Bykova, T. O., Triseleva, T. A., Ivashov, A. V., & Safonkin, A. F. (2016). Morphogenetic Diversity of the Honeybee *Apis mellifera* L. from the Mountain-Forest Zone of Crimea. *Biology Bulletin*, 43(6), 541-546, doi: <https://doi.org/10.1134/S1062359016060054>
- Canal, N. A., Hernández-Ortiz, V., Salas, J. O., & Selivon, D. (2015). Morphometric study of third-instar larvae from five morphotypes of the *Anastrepha fraterculus* cryptic species complex (Diptera, Tephritidae). *ZooKeys*, (540), 41-59, doi: <https://doi.org/10.3897/zookeys.540.6012>
- Carpana, E. (2004). IL Genere *Apis*: Evoluzione e Biogeografia. Parte 1. En *L'aperegina: allevamento e selezione*. (pp. 23-89). Italia: Istituto Nazionale di Apicoltura Bologna
- Carvajal, T. M., Hernandez, L. F., Ho, H. T., Cuenca, M. G., Orantia, B. M., Estrada, C. R., . . . Watanabe, K. (2015). Spatial analysis of wing geometry in dengue vector mosquito, *Aedes aegypti* (L.) (Diptera: Culicidae), populations in Metropolitan Manila, Philippines. *J. Vector Borne Dis*, 53, 127-135. https://pdfs.semanticscholar.org/4c92/372fb188c7441b3f6f327bd3f4b004aacdc2.pdf?_ga=2.10842674.787354217.1581067777-1193939830.1581067777
- Chapman, N. C., Harpur, B. A., Lim, J., Rinderer, T. E., Allsopp, M. H., Zayed, A., & Oldroyd, B. P. (2015). A SNP test to identify Africanized honeybees via proportion of 'African' ancestry. *Molecular Ecology Resources*, 15(6), 1346-1355, doi: <https://doi.org/10.1111/1755-0998.12411>
- Charistos, L., Hatjina, F., Bouga, M., Mladenovic, M., & Maistros, A. D. (2014). Morphological discrimination of Greek honey bee populations based on geometric Morphometrics analysis of wing shape. *Journal of Apicultural Science*, 58(1), 75-84, doi: <https://doi.org/10.2478/jas-2014-0007>
- Chávez-Galarza, J., Garnery, L., Henriques, D., Neves, C. J., Loucif-Ayad, W., Johnston, J. S., & Pinto, A. (2017). Mitochondrial DNA variation of *Apis mellifera iberiensis*: further insights from a large-scale study using sequence data of the tRNA^{Leu-cox2} intergenic region. *Apidologie*, 48(4), 533-544, doi: <https://doi.org/10.1007/s13592-017-0498-2>

- Chávez-Galarza, J., Henriques, D., Johnston, J. S., Carneiro, M., Rufino, J., Patton, J. C., & Pinto, M. A. (2015, Jun). Revisiting the Iberian honey bee (*Apis mellifera iberiensis*) contact zone: maternal and genome-wide nuclear variations provide support for secondary contact from historical refugia. *Mol Ecol.*, 24(12), 2973-92, doi: <https://doi.org/10.1111/mec.13223>
- Combey, R., Quandahor, P., & Mensah, B. A. (2018). Geometric Morphometrics Captures Possible Segregation Occurring within Subspecies *Apis Mellifera Adansonii* in Three Agro Ecological Zones. *Annals of Biological Research*, 9(3), 31-43. Retrieved on February 23, 2019, from: <https://www.scholarsresearchlibrary.com/article/s/geometric-morphometrics-captures-possible-segregation-occurring-within-subspecies-apis-mellifera-adansonii-in-three-agro-ecologica-15155.html>
- Coroian, C. O., Muñoz, I., Schlüns, E. A., Paniti-Teleky, O. R., Erler, S., Furdui, E. M., ..., Moritz, R. F. A. (2014). Climate rather than geography separates two European honeybee subspecies. *Molecular Ecology*, 23(9), 2353-2361. doi: <https://doi.org/10.1111/mec.12731>
- Crozier, R., & Crozier, Y. (1993). The mitochondrial genome of the honeybee *Apis mellifera*: complete sequence and genome organization. *Genetics*, 133, 97-117. Retrieved on March 15, 2019, from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1205303/pdf/ge133197.pdf>
- Da Silva, F., Grassi, M., Sella, M., Francoy, T. M., & Reali, A. H. (2015). Evaluating classification and feature selection techniques for honeybee subspecies identification using wing images. *Computers and Electronics in Agriculture*, 114, 68-77, doi: <https://doi.org/10.1016/j.compag.2015.03.012>
- Daly, H., Hoelmer, K., Norman, P., & Allen, T. (1982). Computer-Assisted Measurement and Identification of Honey Bees (Hymenoptera: Apidae). *Annals of the Entomological Society of America*, 75(6), 591-594, doi: <https://doi.org/10.1093/aesa/75.6.591>
- Darger, K. (2013). *Determining low levels of Africanization in unmanaged honey bee colonies using three diagnostic techniques*. (Master of Science in Entomology), University of Delaware, Estados Unidos. http://udspace.udel.edu/bitstream/handle/19716/12667/Katherine_Darger_thesis.pdf?sequence=1&isAllowed=y
- De Souza, D. A., Wang, Y., Kaftanoglu, O., De Jong, D., Amdam, G. V., Gonçalves, L. S., & Francoy, T. M. (2015). Morphometric identification of queens, workers and intermediates in in vitro reared honey bees (*Apis mellifera*). *PLoS ONE*, 10(4), e0123663, doi: <https://doi.org/10.1371/journal.pone.0123663>
- Dupraw, E. (1965). Non-Linnean taxonomy and the systematics of honeybees. *Systematic Zoology*, 14(1), 1-24, doi: <https://doi.org/10.2307/2411899>
- Eimanifar, A., Kimball, R., Braun, E. L., & Ellis, J. D. (2018). Mitochondrial genome diversity and population structure of two western honey bee subspecies in the Republic of South Africa. *Scientific reports*, 8(1), 1333, doi: <https://doi.org/10.1038/s41598-018-19759-3>
- Eimanifar, A., T. Kimball, R., L. Braun, E., & Ellis, J. D. (2016). The complete mitochondrial genome of the hybrid honey bee, *Apis mellifera capensis* × *Apis mellifera scutellata*, from South Africa. *Mitochondrial DNA Part B*, 1(1), 856-857, doi: <https://doi.org/10.1080/23802359.2016.1250132>
- Francoy, T. M., Wittmann, D., Drauschke, M., Müller, S., Steinhage, V., Bezerra-Laure, M. A. F., . . . Gonçalves, L. (2008). Identification of Africanized honey bees through wing morphometrics: two fast and efficient procedures. *Apidologie*, 39(5), 488-494, doi: <https://doi.org/10.1051/apido:2008028>
- Francoy, T., de Faria Franco, F., & Roubik, D. (2012). Integrated landmark and outline-based morphometric methods efficiently distinguish species of *Euglossa* (Hymenoptera, Apidae, Euglossini). *Apidologie*, 43(6), 609-617, doi: <https://doi.org/10.1007/s13592-012-0132-2>
- Garcia, R., Oliveira, N. T. de, Camargo, S. C., Pires, B., Oliveira, C. de, Teixeira, R., & Pickler, M. (2013). Honey and propolis production, hygiene and defense behaviors of two generations of Africanized honey bees. *Scientia Agricola*, 70(2), 74-81, <https://doi.org/10.1590/S0103-90162013000200003>
- Garnery, L., Solignac, M., Celebrano, G., & Cornuet, J. (1993). A simple test using restricted PCR-amplified mitochondrial DNA to study the genetic structure of *Apis mellifera* L. *Experientia*, 49(11), 1016-1021. doi: <https://doi.org/10.1007/BF02125651>
- Genchi, M. L., Reynaldi, F. J., & Bravi, C. M. (2018). An update of Africanization in honey bee (*Apis mellifera*) populations in Buenos Aires, Argentina. *Journal of Apicultural Research*, 57(5), 611-614. <https://doi.org/10.1080/00218839.2018.1494887>
- Goetze, G. (1940). *Die beste Biene: Züchtungs- und Rassen-Kunde der Honbigbiene nach dem heutigen Stand von Wissenschaft und Praxis*. Germany: Leipzig: Liedloff: Loth und Michaelis.
- Graciano, L. A. (2018). *Niveles de infestación de Varroa destructor (Mesostigmata: Varroidae) en abejas africanizadas (Apis melliferas cutellata)*. (Tesis Magister en Ciencias: Entomología), Universidad Nacional de Colombia-Medellín, Colombia. Retrieved on March 12, 2019, from:

- <http://bdigital.unal.edu.co/64945/1/1128406607.2018.pdf>
- Haddad, N., Adjlane, N., Loucif-Ayad, W., Dash, A., Naganeeswaran, S., Rajashekar, B., ... Sicheritz-Pontén, T. (2017). Mitochondrial genome of the North African Sahara Honeybee, *Apis mellifera sahariensis* (Hymenoptera: Apidae). *Mitochondrial DNA Part B*, 2(2), 548-549, doi: <https://doi.org/10.1080/23802359.2017.1365647>
- Hall, M. J. R., MacLeod, N., & Wardhana, A. H. (2014). Use of wing morphometrics to identify populations of the Old World screw worm fly, *Chrysomya bezziana* (Diptera: Calliphoridae): A preliminary study of the utility of museum specimens. *Acta Tropica*, 138, S49-S55. doi: <https://doi.org/10.1016/j.actatropica.2014.03.023>
- Hamiduzzaman, M., Guzmán-Novoa, E., Goodwin, P. H., Reyes-Quintana, M., Koleoglu, G., Correa-Benítez, A., & Petukhova, T. (2015). Differential responses of Africanized and European honey bees (*Apis mellifera*) to viral replication following mechanical transmission or *Varroa destructor* parasitism. *Journal of invertebrate pathology*, 126, 12-20. doi: <https://doi.org/10.1016/j.jip.2014.12.004>
- Harpur, B. A., Kent, C.F., Molodtsova, D., Lebon, J. M. D., Alqarni, A.S., Owayss, A. A., & Zayed, A. (2014). Population genomics of the honey bee reveals strong signatures of positive selection on worker traits. *Proceedings of the National Academy of Sciences, USA*, 111, 2614-2619, doi: <https://doi.org/10.1073/pnas.1315506111>
- Henriques, D., Parejo, M., Vignal, A., Wragg, D., Wallberg, A., Webster, M. T., & Pinto, M. A. (2018). Developing reduced SNP assays from whole-genome sequence data to estimate introgression in an organism with complex genetic patterns, the Iberian honeybee (*Apis mellifera iberiensis*). *Evolutionary applications*, 11(8), 1270-1282, doi: <https://doi.org/10.1111/eva.12623>
- Hidalgo, M., & Mena, S. (2003). *Proyecto de viabilidad de implementación de una granja apícola en la parroquia de Nanegalito*. (Tesis de Ingeniería en Ciencias Administrativas Tesis de Ingeniería, inédita), Pontificia Universidad Católica del Ecuador, Quito.
- Howell V., Daly, & Balling, S. (1978). Identification of Africanized Honeybees in the Western Hemisphere by Discriminant Analysis. *Journal of the Kansas Entomological Society*, 51(4): 857-869.
- Ivanova, E., Bouga, M., Staykova, T., Mladenovic, M., Rasic, S., Charistos, L., ... Petrov, P. (2012). The genetic variability of honey bees from the Southern Balkan Peninsula, based on alloenzymic data. *Journal of Apicultural Research*, 51(4), 329-335, doi: <https://doi.org/10.3896/IBRA.1.51.4.06>
- Jarnevic, C. S., Esaias, W. E., Ma, P. L. A., Morissette, J. T., Nickeson, J. E., Stohlgren, T. J., ... Tan, B. (2014). Regional distribution models with lack of proximate predictors: Africanized honeybees expanding north. *Diversity and Distributions*, 20(2), 193-201, doi: <https://doi.org/10.1111/ddi.12143>
- Kandemir, İ., Özkan, A., & Fuchs, S. (2011). Reevaluation of honeybee (*Apis mellifera*) microtaxonomy: a geometric morphometric approach. *Apidologie*, 42(5), 618, doi: <https://doi.org/10.1007/s13592-011-0063-3>
- Kelomey, A., Paraíso, A. A., Sina, H., Legout, H., Adjanohoun, A., Garnery, L., & Baba-Moussa, L. (2017). Genetic Variability of the Mitochondrial DNA in Honeybees (*Apis mellifera* L.) from Benin. *Journal of Agricultural Science and Technology*, 7, 557-566, doi: <https://doi.org/10.17265/2161-6256/2017.08.006>
- Kerr, W. E. (1967). The history of the introduction of African bees in Brazil. *South African Bee J.*, 39: 33-35.
- Kono, Y., & Kohn, J. (2015). Range and frequency of africanized honey bees in California (USA). *PLoS ONE*, 10(9), e0137407, doi: <https://doi.org/10.1371/journal.pone.0137407>
- Magnus, R., & Szalanski, A. L. (2010). Genetic evidence for honey bees (*Apis mellifera* L.) of Middle Eastern lineage in the United States. *Sociobiology*, 55(1B), 285-296. Retrieved on March 15, 2019, from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.630.4651&rep=rep1&type=pdf>
- Medina-Flores, C. A., Guzmán-Novoa, E., Hamiduzzaman, M. M., Aguilera Soto, J., Carlos, L., & Marco, A. (2015). Africanización de colonias de abejas melíferas (*Apis mellifera*) en tres regiones climáticas del norte de México. *Veterinaria México OA*, 2(4), 1-9. Retrieved on March 2, 2019, from: <https://www.redalyc.org/pdf/4935/493548765001.pdf>
- Meixner, M. D., Pinto, M. A., Bouga, M., Kryger, P., Ivanova, E., & Fuchs, S. (2013). Standard methods for characterising subspecies and ecotypes of *Apis mellifera*. *Journal of Apicultural Research*, 52(4), 1-28, doi: <https://doi.org/10.3896/IBRA.1.52.4.05>
- Mendoza, Y., Antúnez, K., Branchiccela, B., Anido, M., Santos, E., & Invernizzi, C. (2014). Nosemaceranae and RNA viruses in European and Africanized honeybee colonies (*Apis mellifera*) in Uruguay. *Apidologie*, 45(2), 224-234, doi: <https://doi.org/10.1007/s13592-013-0241-6>
- Miguel, I., Garnery, L., Iriondo, M., Baylac, M., Manzano, C., Steve, W., & Estonba, A. (2016).

- Origin, evolution and conservation of the honey bees from La Palma Island (Canary Islands): molecular and morphological data. *Journal of Apicultural Research*, 54(5), 427-440, doi: <https://doi.org/10.1080/00218839.2016.1180017>
- Moore, P., Wilson, M., & Skinner, J. (2015). *Africanized Bees: Better Understanding, Better Prepared*. USA: University of Tennessee, Department of Entomology and Plant Pathology, the, Knoxville TN. Retrieved on February 16, 2019, from: <https://bee-health.extension.org/africanized-bees-better-understanding-better-prepared/>
- Mortensen, A., & Ellis, J. (2015). The frequency of African (*Apis mellifera scutellata* Lepeletier) matrilineal usurpation of managed European-derived honey bee (*A. mellifera* L.) colonies in the southeastern United States. *Insectes Sociaux*, 62(2), 151-155. doi: <https://doi.org/10.1007/s00040-014-0383-1>
- Muñoz, I., Enriques, D., Jara, L., Johnston, J.S., Chávez-Galarza, J., Rua, P. de la, & Pinto, M. (2016). SNPs selected by information content outperform randomly selected microsatellite loci for delineating genetic identification and introgression in the endangered dark European honeybee (*Apis mellifera mellifera*). *Mol. Ecol. Res.*, 17(4), 783-795, doi: <https://doi.org/10.1111/1755-0998.12637>
- Muñoz, I., Henriques, D., Johnston, S., Chávez-Galarza, J., Kryger, P., & Pinto, M. A. (2015). Reduced SNP Panels for Genetic Identification and Introgression Analysis in the Dark Honey Bee (*Apis mellifera mellifera*). *PLoS ONE*, 10(4), e0124365, doi: <https://doi.org/10.1371/journal.pone.0124365>
- Muñoz, I., Lodesani, M., Rúa, P. de la, & Dall'Olio, R. (2014). Estimating introgression in *Apis mellifera siciliana* populations: are the conservation islands really effective? *Insect Conserv Divers*, 7(6), 563-571, doi: <https://doi.org/10.1111/icad.12092>
- Nogueira-Neto, P. (1972). Notas sobre a história da apicultura brasileira. En J. M. F. De Camargo (ed.), *Manual de Apicultura*. (pp. 17-32). Sao Paulo, Brasil: Editora Agronomia Ceres.
- Nunamaker, R. A., & Wilson, W. T. (1982). Isozyme changes in the honeybee, *Apis mellifera* L., during larval morphogenesis. *Insect Biochemistry*, 12(1), 99-104, doi: [https://doi.org/10.1016/0020-1790\(82\)90076-2](https://doi.org/10.1016/0020-1790(82)90076-2)
- Nunes, L. A., Araújo, E. D. de, Marchini, L. C., & Moreti, A. C. de C. C. (2012). Variation morphogeometrics of Africanized honey bees (*Apis mellifera*) in Brazil. *Iheringia. Série Zoologia*, 102(3), 321-326, doi: <http://dx.doi.org/10.1590/S0073-47212012005000002>
- Oleksa, A., & Tofilski, A. (2015). Wing geometric morphometrics and microsatellite analysis provide similar discrimination of honey bee subspecies. *Apidologie*, 46(1), 49-60, doi: <https://doi.org/10.1007/s13592-014-0300-7>
- Ostroverkhova, N. V., Konusova, O. L., Kucher, A. N., Kireeva, T. N., Vorotov, A. A., & Belikh, E. A. (2015). Genetic diversity of the locus COI-COII of mitochondrial DNA in honeybee populations (*Apis mellifera* L.) from the Tomsk region. *Russian Journal of Genetics*, 51(1), 80-90, doi: <https://doi.org/10.1134/S102279541501010X>
- Oyerinde, A., Salako, E., & Rabiou, M. (2017). Morphometric taxonomy of honeybee races of *Apis mellifera* L. in Kaduna state. *Journal of Entomology and Zoology Studies*, 5(5), 825-829. Retrieved on March 23, 2019, from: <http://www.entomoljournal.com/archives/2017/vol5issue5/PartK/4-3-187-692.pdf>
- Péntek-Zakar, E., Oleksa, A., Borowik, T., & Kusza, S. (2015). Population structure of honey bees in the Carpathian Basin (Hungary) confirms introgression from surrounding subspecies *Ecology and Evolution*, 5(23), 5456-5467, doi: <https://doi.org/10.1002/ece3.1781>
- Pinto, M. A., Henriques, D., Chavez-Galarza, J., Kryger, P., Garnery, L., van der Zee, R.,... Johnston, J. S. (2014). Genetic integrity of the Dark European honey bee (*Apis mellifera mellifera*) from protected populations: a genome-wide assessment using SNPs and medina sequence data. *Journal of Apicultural Research*, 53(2), 269-278, doi: <https://doi.org/10.3896/IBRA.1.53.2.08>
- Portman, Z. M., Tepedino, V. J., Tripodi, A. D., Szalanski, A. L., & Durham, S. L. (2018). Local extinction of a rare plant pollinator in Southern Utah (USA) associated with invasion by Africanized honey bees. *Biological Invasions*, 20(3), 593-606, doi: <https://doi.org/10.1007/s10530-017-1559-1>
- Qubaiová, J., Růžička, J., & Šípková, H. (2015). Taxonomic revision of genus *Ablattaria* Reitter (Coleoptera, Silphidae) using geometric morphometrics. *ZooKeys*, (477), 79-142, doi: <https://doi.org/10.3897/zookeys.477.8446>
- Rangel, J., Giresi, M., Pinto, M. A., Baum, K. A., Rubink, W. L., Coulson, R. N., & Johnston, J. S. (2016). Africanization of a feral honey bee (*Apis mellifera*) population in South Texas: does a decade make a difference? *Ecology and Evolution*, 6(7), 2158-2169, doi: <https://doi.org/10.1002/ece3.1974>
- Rohlf, F. J., & Marcus, L. F. (1993). A revolution morphometrics. *Trends in Ecology & Evolution*, 8(4), 129-132. doi: [https://doi.org/10.1016/0169-5347\(93\)90024-J](https://doi.org/10.1016/0169-5347(93)90024-J)
- Rortais, A., Arnold, G., Alburaki, M., Legout, H., & Garnery, L. (2011). Review of the Dral COI-COII test for the conservation of the black honeybee (*Apis mellifera mellifera*). *Conserv Genet Resour*, 3(2), 383-391, doi: <https://doi.org/10.1007/s12686-010-9351-x>

- Rúa, P. de la, Jaffé, R., Dall'Olio, R., Muñoz, I., & Serrano, J. (2009). Biodiversity, conservation and current threats to European honeybees. *Apidologie*, 40(3), 263-284, doi: <https://doi.org/10.1051/apido/2009027>
- Rúa, P. de la, Martínez, J., Domingo, O., & Gabaldón, I. (2013). Caracterización molecular de la biodiversidad de la cabaña apícola de la provincia de Albacete. *Revista de Estudios Albacetenses*, (9), 175-196. Retrieved on April 6, 2019, from: <https://dialnet.unirioja.es/descarga/articulo/4252286.pdf>
- Ruttner, F. (1988). *Biogeography and Taxonomy of Honeybees*. Springer, Berlin, doi: <http://dx.doi.org/10.1007/978-3-642-72649-1>
- Ruttner, F. (1992). Natural history of honey bees. Munich, Germany: Ehrenwirth Verlag.
- Ruttner, F., Tassencourt, L., & Louveaux, J. (1978). Biometrical-statistical analysis of the geographic variability of *Apis mellifera* L. Material and methods. *Apidologie*, 9(4), 363-381, doi: <https://doi.org/10.1051/apido:19780408>
- Sanford, M. (2006). Africanized honey bee: A biological revolution with human cultural implications. *American Bee Journal*. Retrieved on February 5, 2019, from: http://apisenterprises.com/papers_hm/Misc/AH/B%20in%20the%20Americas.htm
- Santana, F. S., Costa, A. H., Truzzi, F. S., Silva, F. L., Santos, S. L., Franco, T. M., & Saraiva, A. M. (2014). A reference process for automating bee species identification based on wing images and digital image processing. *Ecological Informatics*, 24, 248-260, doi: <https://doi.org/10.1016/j.ecoinf.2013.12.001>
- Sousa, A. R. S., Araújo, E. D., Gramacho, K. P., & Nunes, L. A. (2016). Bee's morphometrics and behavior in response to seasonal effects from ecoregions. *Genetics and Molecular Research*, 15(2), 1-14, doi: <http://dx.doi.org/10.4238/gmr.150275>
- Strauss, U., Dietemann, V., Human, H., Crewe, R. M., & Pirk, C. W. (2015). Resistance rather than tolerance explains survival of savannah honeybees (*Apis mellifera* scutellata) to infestation by the parasitic mite *Varroa destructor*. *Parasitology*, 143(3), 374-387, doi: <https://doi.org/10.1017/S0031182015001754>
- Su, X., Cai, X., & Huang, D. (2018). Taxonomic Ecology of Geometric Morphometry on Classification and Identification of Sphingid Moths (Lepidoptera: Sphingidae). *Ekoloji*, 27(106), 827-835. Retrieved on March 15, 2019, from: <http://www.ekolojidergisi.com/download/taxonomic-ecology-of-geometric-morphometry-on-classification-and-identification-of-sphingid-moths-5466.pdf>
- Sylvester, H. A., & Rinderer, T. E. (1987). Fast Africanized Bee Identification System (FABIS). *American Bee Journal*, 127(7), 511-516.
- Szalanski, A. L., & Magnus, R. M. (2010). Mitochondrial DNA characterization of Africanized honey bee (*Apis mellifera* L.) populations from the USA. *Journal of Apicultural Research*, 49(2), 177-185, doi: <https://doi.org/10.3896/IBRA.1.49.2.06>
- Szalanski, A., & Tripodi, A. (2014). Assessing the Utility of a PCR Diagnostics Marker for the Identification of Africanized Honey Bee, *Apis mellifera* L., (Hymenoptera: Apidae) in the United States. *Sociobiology*, 61(2), 234-236, doi: <http://dx.doi.org/10.13102/sociobiology.v61i2.234-236>
- Takahashi, J.-i., Wakamiya, T., Kiyoshi, T., Uchiyama, H., Yajima, S., Kimura, K., & Nomura, T. (2016). The complete mitochondrial genome of the Japanese honeybee, *Apis cerana japonica* (Insecta: Hymenoptera: Apidae). *Mitochondrial DNA Part B*, 1(1), 156-157, doi: <https://doi.org/10.1080/23802359.2016.1144108>
- Techer, M. A., Clémencet, J., Turpin, P., Volbert, N., Reynaud, B., & Delatte, H. (2015). Genetic characterization of the honeybee (*Apis mellifera*) population of Rodrigues Island, based on microsatellite and mitochondrial DNA. *Apidologie*, 46(4), 445-454, doi: <https://doi.org/10.1007/s13592-014-0335-9>
- Techer, M., Clémencet, J., Simiand, C., Preaduth, S., Azali, H. A., Reynaud, B., & Hélène, D. (2017). Large-scale mitochondrial DNA analysis of native honey bee *Apis mellifera* populations reveals a new African subgroup private to the South West Indian Ocean islands. *BMC genetics*, 18(1), 53, doi: <https://doi.org/10.1186/s12863-017-0520-8>
- Tibatá, V. M., Arias, E., Corona, M., Ariza Botero, F., Figueroa-Ramírez, J., & Junca, H. (2018). Determination of the Africanized mitotypes in populations of honey bees (*Apis mellifera* L.) of Colombia. *Journal of Apicultural Research*, 57(2), 219-227, doi: <https://doi.org/10.1080/00218839.2017.1409065>
- Torcida, S., & Perez, I. S. (2012). Análisis de Procrustes y el estudio de la variación morfológica. *Revista Argentina de Antropología Biológica*, 14(1), 131-141. Retrieved on March 15, 2019, from: <https://revistas.unlp.edu.ar/raab/article/view/537>
- Valido, A., Rodríguez-Rodríguez, M. C., & Jordano, P. (2014). Impacto de la introducción de la abeja doméstica (*Apis mellifera*, Apidae) en el Parque Nacional del Teide (Tenerife, Islas Canarias). *Revista Ecosistemas*, 23(3), 58-66, doi: <https://doi.org/10.7818/ECOS.2014.23-3.08>
- Verde, M. M. (2014). Apicultura y seguridad alimentaria. *Revista Cubana de Ciencia*

- Agrícola*, 48(1), 25-31. Retrieved on March 23, 2019, from:
<https://www.redalyc.org/pdf/1930/193030122008.pdf>
- Wallberg, A., Han, F., Wellhagen, G., Dahle, B., Kawata, M., Haddad, N., . . . Rúa, P. de la (2014). A worldwide survey of genome sequence variation provides insight into the evolutionary history of the honeybee *Apis mellifera*. *Nature Genetics*, 46(10), 1081-1088, doi: <https://doi.org/10.1038/ng.3077>
- Whitfield, C., Behura, S., Berlocher, S., Clark, A., Johnston, S., Sheppard, W., . . . Tsutsui, N. (2006). Thrice out of Africa: ancient and recent expansions of the honey bee, *Apis mellifera*. *Science*, 314(5799), 642-645, doi: <https://doi.org/10.1126/science.1132772>
- Winston, M. (1992). Killer bees. The Africanized honey bee in the Americas. Cambridge, Massachusetts, USA: Harvard University Press.
- Wu, M.-C., Lu, T.-H., & Lu, K.-H. (2017). PCR-RFLP of mitochondrial DNA reveals two origins of *Apis mellifera* in Taiwan. *Saudi Journal of Biological Sciences*, 24(5), 1069-1074. doi: <https://doi.org/10.1016/j.sjbs.2016.12.008>
- Zelditch, M., Swiderski, D., Sheets, D., & Fink, W. L. (2004). Geometric Morphometrics for Biologists: A Primer: Academic Press.
- Zhao, W., Tan, K., Zhou, D., Wang, M., Cheng, C., Yu, Z., . . . He, S. (2014). Phylogeographic analysis of *Apis cerana* populations on Hainan Island and southern mainland China, based on mitochondrial DNA sequences. *Apidologie*, 45(1), 21-33, doi: <https://doi.org/10.1007/s13592-013-0223-8>